



# LAGRANGE DESCRIPTION III EULER DESCRIPTION I

**Jiří Plešek**

Institute of Thermomechanics  
Czech Academy of Sciences



# Contents

---

- Lagrange description
  - Polar decomposition
  - A general strain tensor
- Euler description
  - Trajectory
  - Velocity
  - Acceleration



# Rotation

---

Special case

$$[F] = \text{orthonormal} = [R]$$



# Rotation

---

Special case

$$[F] = \text{orthonormal} = [R]$$

GL strain tensor

$$[e] = \frac{1}{2}([F]^T[F] - [I]) = \frac{1}{2}([R]^T[R] - [I]) = [0]$$



# Rotation

---

Special case

$$[F] = \text{orthonormal} = [R]$$

GL strain tensor

$$[e] = \frac{1}{2}([F]^T[F] - [I]) = \frac{1}{2}([R]^T[R] - [I]) = [0]$$





# Stretching

---

Special case

$$[F] = \text{sym} + \text{def} = [U]$$





# Stretching

---

Special case

$$[F] = \text{sym} + \text{def} = [U]$$

Principal axes

$$[U] \text{ sym} \mapsto [U'] = [\Lambda] = \text{diag}[\lambda_1, \lambda_2, \lambda_3] \quad (\text{principal stretches} > 0)$$



# Stretching

---

Special case

$$[F] = \text{sym} + \text{def} = [U]$$

Principal axes

$$[U] \text{ sym} \mapsto [U'] = [\Lambda] = \text{diag}[\lambda_1, \lambda_2, \lambda_3] \quad (\text{principal stretches} > 0)$$

Line segment

$$d\{x'\} = [F'] d\{X'\} = [U'] d\{X'\} = [\Lambda] d\{X'\}$$





# Stretching

---

Special case

$$[F] = \text{sym} + \text{def} = [U]$$

Principal axes

$$[U] \text{ sym} \mapsto [U'] = [\Lambda] = \text{diag}[\lambda_1, \lambda_2, \lambda_3] \quad (\text{principal stretches} > 0)$$

Line segment

$$d\{x'\} = [F'] d\{X'\} = [U'] d\{X'\} = [\Lambda] d\{X'\}$$

$$\left. \begin{aligned} dx'_1 &= \lambda_1 dX'_1 \\ dx'_2 &= \lambda_2 dX'_2 \\ dx'_3 &= \lambda_3 dX'_3 \end{aligned} \right\} \begin{array}{l} \lambda_k > 1 \text{ tension} \\ \lambda_k < 1 \text{ compression} \end{array}$$



# Stretching

---

Special case

$$[F] = \text{sym} + \text{def} = [U]$$

Principal axes

$$[U] \text{ sym} \mapsto [U'] = [\Lambda] = \text{diag}[\lambda_1, \lambda_2, \lambda_3] \quad (\text{principal stretches} > 0)$$

Line segment

$$d\{x'\} = [F'] d\{X'\} = [U'] d\{X'\} = [\Lambda] d\{X'\}$$

$$\left. \begin{aligned} dx'_1 &= \lambda_1 dX'_1 \\ dx'_2 &= \lambda_2 dX'_2 \\ dx'_3 &= \lambda_3 dX'_3 \end{aligned} \right\} \begin{array}{l} \lambda_k > 1 \text{ tension} \\ \lambda_k < 1 \text{ compression} \end{array}$$





# Composition

---

Stretch and rotation

$$d\{x_m\} = [U] d\{X\} \text{ followed by } d\{x\} = [R] d\{x_m\}$$



# Composition

---

Stretch and rotation

$$d\{x_m\} = [U] d\{X\} \text{ followed by } d\{x\} = [R] d\{x_m\} \Rightarrow d\{x\} = [R][U] d\{X\}$$



# Composition

---

Stretch and rotation

$$d\{x_m\} = [U] d\{X\} \text{ followed by } d\{x\} = [R] d\{x_m\} \Rightarrow d\{x\} = [R][U] d\{X\}$$

Hence

$$[F] = [R][U]$$



# Composition

---

Stretch and rotation

$$d\{x_m\} = [U] d\{X\} \text{ followed by } d\{x\} = [R] d\{x_m\} \Rightarrow d\{x\} = [R][U] d\{X\}$$

Hence

$$\mathbf{F} = \mathbf{R}\mathbf{U}$$





# Composition

---

Stretch and rotation

$$d\{x_m\} = [U] d\{X\} \text{ followed by } d\{x\} = [R] d\{x_m\} \Rightarrow d\{x\} = [R][U] d\{X\}$$

Hence

$$\mathbf{F} = \mathbf{R}\mathbf{U}$$

Multiplicative decomposition

$$\mathbf{F} = \mathbf{F}_N \mathbf{F}_{N-1} \cdots \mathbf{F}_2 \mathbf{F}_1$$



# Composition

---

Stretch and rotation

$$d\{x_m\} = [U] d\{X\} \text{ followed by } d\{x\} = [R] d\{x_m\} \Rightarrow d\{x\} = [R][U] d\{X\}$$

Hence

$$\mathbf{F} = \mathbf{R}\mathbf{U}$$

Multiplicative decomposition

$$\mathbf{F} = \mathbf{F}_N \mathbf{F}_{N-1} \cdots \mathbf{F}_2 \mathbf{F}_1$$

Rotation followed by stretch

$$\mathbf{F} = \mathbf{R}\mathbf{U} = \mathbf{R}\mathbf{U}\mathbf{R}^T \mathbf{R}$$



# Composition

---

Stretch and rotation

$$d\{x_m\} = [U] d\{X\} \text{ followed by } d\{x\} = [R] d\{x_m\} \Rightarrow d\{x\} = [R][U] d\{X\}$$

Hence

$$\mathbf{F} = \mathbf{R}\mathbf{U}$$

Multiplicative decomposition

$$\mathbf{F} = \mathbf{F}_N \mathbf{F}_{N-1} \cdots \mathbf{F}_2 \mathbf{F}_1$$

Rotation followed by stretch

$$\mathbf{F} = \mathbf{R}\mathbf{U} = \mathbf{R}\mathbf{U}\mathbf{R}^T\mathbf{R} = \mathbf{V}\mathbf{R} \quad (\text{same principal stretches})$$



# Decomposition

---

$$\text{If } \mathbf{F} = \mathbf{R}\mathbf{U} \text{ then } \mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}} \text{ sym+def}$$



# Decomposition

---

If  $\mathbf{F} = \mathbf{R}\mathbf{U}$  then  $\mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}}$  sym+def

Set  $\mathbf{R} = \mathbf{F}\mathbf{U}^{-1}$  and verify  $\mathbf{R}^T\mathbf{R} = \mathbf{U}^{-1}\mathbf{F}^T\mathbf{F}\mathbf{U}^{-1} = \mathbf{U}^{-1}\mathbf{U}^2\mathbf{U}^{-1} = \mathbf{I}$



# Decomposition

---

If  $\mathbf{F} = \mathbf{R}\mathbf{U}$  then  $\mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}}$  sym+def

Set  $\mathbf{R} = \mathbf{F}\mathbf{U}^{-1}$  and verify  $\mathbf{R}^T\mathbf{R} = \mathbf{U}^{-1}\mathbf{F}^T\mathbf{F}\mathbf{U}^{-1} = \mathbf{U}^{-1}\mathbf{U}^2\mathbf{U}^{-1} = \mathbf{I}$

... which proves existence of  $\mathbf{F} = \mathbf{R}\mathbf{U}$  for any regular  $\mathbf{F}$ .

---





# Decomposition

---

If  $\mathbf{F} = \mathbf{R}\mathbf{U}$  then  $\mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}}$  sym+def

Set  $\mathbf{R} = \mathbf{F}\mathbf{U}^{-1}$  and verify  $\mathbf{R}^T\mathbf{R} = \mathbf{U}^{-1}\mathbf{F}^T\mathbf{F}\mathbf{U}^{-1} = \mathbf{U}^{-1}\mathbf{U}^2\mathbf{U}^{-1} = \mathbf{I}$

... which proves existence of  $\mathbf{F} = \mathbf{R}\mathbf{U}$  for any regular  $\mathbf{F}$ .

---

Square root of a symmetric, positive definite matrix:

1.  $[C] = [F]^T[F]$  sym+def for any regular  $[F]$  (Cauchy-Green metric tensor)

# Decomposition

---

If  $\mathbf{F} = \mathbf{R}\mathbf{U}$  then  $\mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}}$  sym+def

Set  $\mathbf{R} = \mathbf{F}\mathbf{U}^{-1}$  and verify  $\mathbf{R}^T\mathbf{R} = \mathbf{U}^{-1}\mathbf{F}^T\mathbf{F}\mathbf{U}^{-1} = \mathbf{U}^{-1}\mathbf{U}^2\mathbf{U}^{-1} = \mathbf{I}$

... which proves existence of  $\mathbf{F} = \mathbf{R}\mathbf{U}$  for any regular  $\mathbf{F}$ .

---

Square root of a symmetric, positive definite matrix:

1.  $[C] = [F]^T[F]$  sym+def for any regular  $[F]$  (Cauchy-Green metric tensor)
2. The eigenvalue problem:  $[C]\{\varphi_k\} = \mu_k\{\varphi_k\}$  ,  $\mu_k > 0$



# Decomposition

---

If  $\mathbf{F} = \mathbf{R}\mathbf{U}$  then  $\mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}}$  sym+def

Set  $\mathbf{R} = \mathbf{F}\mathbf{U}^{-1}$  and verify  $\mathbf{R}^T\mathbf{R} = \mathbf{U}^{-1}\mathbf{F}^T\mathbf{F}\mathbf{U}^{-1} = \mathbf{U}^{-1}\mathbf{U}^2\mathbf{U}^{-1} = \mathbf{I}$

... which proves existence of  $\mathbf{F} = \mathbf{R}\mathbf{U}$  for any regular  $\mathbf{F}$ .

---

Square root of a symmetric, positive definite matrix:

1.  $[C] = [F]^T[F]$  sym+def for any regular  $[F]$  (Cauchy-Green metric tensor)
2. The eigenvalue problem:  $[C]\{\varphi_k\} = \mu_k\{\varphi_k\}$  ,  $\mu_k > 0$
3. Let  $\lambda_k = \sqrt{\mu_k} > 0$  ,  $[\Lambda] = \text{diag}[\lambda_1 \lambda_2 \lambda_3] \Rightarrow [\Lambda]^2 = \text{diag}[\mu_1 \mu_2 \mu_3]$

# Decomposition

---

If  $\mathbf{F} = \mathbf{R}\mathbf{U}$  then  $\mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}}$  sym+def

Set  $\mathbf{R} = \mathbf{F}\mathbf{U}^{-1}$  and verify  $\mathbf{R}^T\mathbf{R} = \mathbf{U}^{-1}\mathbf{F}^T\mathbf{F}\mathbf{U}^{-1} = \mathbf{U}^{-1}\mathbf{U}^2\mathbf{U}^{-1} = \mathbf{I}$

... which proves existence of  $\mathbf{F} = \mathbf{R}\mathbf{U}$  for any regular  $\mathbf{F}$ .

---

Square root of a symmetric, positive definite matrix:

1.  $[C] = [F]^T[F]$  sym+def for any regular  $[F]$  (Cauchy-Green metric tensor)
2. The eigenvalue problem:  $[C]\{\varphi_k\} = \mu_k\{\varphi_k\}$  ,  $\mu_k > 0$
3. Let  $\lambda_k = \sqrt{\mu_k} > 0$  ,  $[\Lambda] = \text{diag}[\lambda_1 \lambda_2 \lambda_3] \Rightarrow [\Lambda]^2 = \text{diag}[\mu_1 \mu_2 \mu_3]$
4. Let  $[U] = [\Phi][\Lambda][\Phi]^T$  sym+def

# Decomposition

---

If  $\mathbf{F} = \mathbf{R}\mathbf{U}$  then  $\mathbf{F}^T\mathbf{F} = \mathbf{U}^T\mathbf{R}^T\mathbf{R}\mathbf{U} = \mathbf{U}\mathbf{U} = \mathbf{U}^2 \Rightarrow \mathbf{U} = \sqrt{\mathbf{F}^T\mathbf{F}}$  sym+def

Set  $\mathbf{R} = \mathbf{F}\mathbf{U}^{-1}$  and verify  $\mathbf{R}^T\mathbf{R} = \mathbf{U}^{-1}\mathbf{F}^T\mathbf{F}\mathbf{U}^{-1} = \mathbf{U}^{-1}\mathbf{U}^2\mathbf{U}^{-1} = \mathbf{I}$

... which proves existence of  $\mathbf{F} = \mathbf{R}\mathbf{U}$  for any regular  $\mathbf{F}$ .

---

Square root of a symmetric, positive definite matrix:

1.  $[C] = [F]^T[F]$  sym+def for any regular  $[F]$  (Cauchy-Green metric tensor)
2. The eigenvalue problem:  $[C]\{\varphi_k\} = \mu_k\{\varphi_k\}$ ,  $\mu_k > 0$
3. Let  $\lambda_k = \sqrt{\mu_k} > 0$ ,  $[\Lambda] = \text{diag}[\lambda_1 \lambda_2 \lambda_3] \Rightarrow [\Lambda]^2 = \text{diag}[\mu_1 \mu_2 \mu_3]$
4. Let  $[U] = [\Phi][\Lambda][\Phi]^T$  sym+def

It follows that  $[U]^2 = [\Phi][\Lambda][\Phi]^T[\Phi][\Lambda][\Phi]^T = [\Phi][\Lambda]^2[\Phi]^T = [C]$ .





# Polar decomposition

---

Theorem

$$\mathbf{F} = \mathbf{R}\mathbf{U} = \mathbf{V}\mathbf{R} \quad \text{uniquely}$$

The right and left stretch tensors

$\mathbf{U}$ ,  $\mathbf{V}$  sym+def, having the same principal stretches  $\lambda_k > 0$

Rotation tensor

$\mathbf{R}$  = proper orthogonal, that is,  $\det |\mathbf{R}| = +1$





# Polar decomposition

---

Theorem

$$\mathbf{F} = \mathbf{R}\mathbf{U} = \mathbf{V}\mathbf{R} \quad \text{uniquely}$$

The right and left stretch tensors

$\mathbf{U}$ ,  $\mathbf{V}$  sym+def, having the same principal stretches  $\lambda_k > 0$

Rotation tensor

$\mathbf{R}$  = proper orthogonal, that is,  $\det |\mathbf{R}| = +1$

...verifies as


$$\det |\mathbf{F}| = \underbrace{\det |\mathbf{R}|}_{\pm 1} \underbrace{\det |\mathbf{U}|}_{> 0} = J > 0$$



# Two families of strain tensors

---

Grand scheme



Lagrange family

$$\mathbf{E} = f(\mathbf{U}) \quad (\text{e.g. Green-Lagrange})$$

Euler family

$$\mathbf{A} = f(\mathbf{V}) \quad (\text{e.g. Euler-Almansi})$$



# Examples of Lagrangean tensors

---

Green-Lagrange

$$\mathbf{e} = \frac{1}{2}(\mathbf{F}^T \mathbf{F} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^T \mathbf{R}^T \mathbf{R} \mathbf{U} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^2 - \mathbf{I})$$

...illogical formula but easy computation.



# Examples of Lagrangean tensors

---

## Green-Lagrange

$$\mathbf{e} = \frac{1}{2}(\mathbf{F}^T \mathbf{F} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^T \mathbf{R}^T \mathbf{R} \mathbf{U} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^2 - \mathbf{I})$$

...illogical formula but easy computation.

## Small strain

$$\boldsymbol{\epsilon} = \frac{1}{2}(\mathbf{z} + \mathbf{z}^T) = \frac{1}{2}(\mathbf{F} + \mathbf{F}^T) - \mathbf{I} = \frac{1}{2}(\mathbf{R} \mathbf{U} + \mathbf{U} \mathbf{R}^T) - \mathbf{I}$$



# Examples of Lagrangean tensors

---

## Green-Lagrange

$$\mathbf{e} = \frac{1}{2}(\mathbf{F}^T \mathbf{F} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^T \mathbf{R}^T \mathbf{R} \mathbf{U} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^2 - \mathbf{I})$$

...illogical formula but easy computation.

## Small strain

$$\boldsymbol{\epsilon} = \frac{1}{2}(\mathbf{z} + \mathbf{z}^T) = \frac{1}{2}(\mathbf{F} + \mathbf{F}^T) - \mathbf{I} = \frac{1}{2}(\mathbf{R} \mathbf{U} + \mathbf{U} \mathbf{R}^T) - \mathbf{I}$$

...depends on rotation but ...

$$\mathbf{R} \rightarrow \mathbf{I}: \quad \boldsymbol{\epsilon} \simeq \mathbf{U} - \mathbf{I} \quad (\text{relative elongation})$$





# Examples of Lagrangean tensors

---

## Green-Lagrange

$$\mathbf{e} = \frac{1}{2}(\mathbf{F}^T \mathbf{F} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^T \mathbf{R}^T \mathbf{R} \mathbf{U} - \mathbf{I}) = \frac{1}{2}(\mathbf{U}^2 - \mathbf{I})$$

...illogical formula but easy computation.

## Small strain

$$\boldsymbol{\epsilon} = \frac{1}{2}(\mathbf{z} + \mathbf{z}^T) = \frac{1}{2}(\mathbf{F} + \mathbf{F}^T) - \mathbf{I} = \frac{1}{2}(\mathbf{R} \mathbf{U} + \mathbf{U} \mathbf{R}^T) - \mathbf{I}$$

...depends on rotation but ...


$$\mathbf{R} \rightarrow \mathbf{I}: \quad \boldsymbol{\epsilon} \simeq \mathbf{U} - \mathbf{I} \quad (\text{relative elongation})$$

## Biot

$$\mathbf{b} = \mathbf{U} - \mathbf{I} \quad \text{for any } \mathbf{R}$$



# EULER DESCRIPTION



$$\vec{N}(\vec{x}, t) = \underbrace{\vec{N}[\vec{x}(\vec{X}, t), t]}_{\text{identical functions}} \equiv \vec{V}(\vec{X}, t)$$



# Composed mappings

---

Composition of a mapping

$$v \circ x : \mathbb{R} \rightarrow \mathbb{R}$$



# Composed mappings

---

Composition of a mapping

$$v \circ x : \mathbb{R} \rightarrow \mathbb{R} \quad \dots \text{ means } V(X) = v(x(X))$$



# Composed mappings

---

Composition of a mapping

$$v \circ x : \mathbb{R} \rightarrow \mathbb{R} \quad \dots \text{ means } V(X) = v(x(X))$$

Example

$$v(x) = \sin x, \quad x = X^2 \quad \Rightarrow \quad v = \sin X^2$$



# Composed mappings

---

Composition of a mapping

$$v \circ x : \mathbb{R} \rightarrow \mathbb{R} \quad \dots \text{means} \quad V(X) = v(x(X))$$

Example

$$v(x) = \sin x, \quad x = X^2 \quad \Rightarrow \quad v = \sin X^2$$

...master caution!

$$\sin X^2 \equiv V(X) \not\equiv v(X) \equiv \sin X$$



# Composed mappings

---

Composition of a mapping

$$v \circ x : \mathbb{R} \rightarrow \mathbb{R} \quad \dots \text{means} \quad V(X) = v(x(X))$$

Example

$$v(x) = \sin x, \quad x = X^2 \quad \Rightarrow \quad v = \sin X^2$$

...master caution!

$$\sin X^2 \equiv V(X) \not\equiv v(X) \equiv \sin X \quad \text{but} \quad V(X) = v(x) \quad \text{for all} \quad x = X^2$$





# Composed mappings

---

Composition of a mapping

$$v \circ x : \mathbb{R} \rightarrow \mathbb{R} \quad \dots \text{means} \quad V(X) = v(x(X))$$

Example

$$v(x) = \sin x, \quad x = X^2 \quad \Rightarrow \quad v = \sin X^2$$

...master caution!

$$\sin X^2 \equiv V(X) \not\equiv v(X) \equiv \sin X \quad \text{but} \quad V(X) = v(x) \quad \text{for all} \quad x = X^2$$

Quiz

$$\mathbf{V}(\mathbf{X}, t) = \mathbf{V}(\mathbf{X}(\mathbf{x}, t), t) = \mathbf{v}(\mathbf{x}, t)$$



# Composed mappings

---

Composition of a mapping

$$v \circ x : \mathbb{R} \rightarrow \mathbb{R} \quad \dots \text{means} \quad V(X) = v(x(X))$$

Example

$$v(x) = \sin x, \quad x = X^2 \quad \Rightarrow \quad v = \sin X^2$$


...master caution!

$$\sin X^2 \equiv V(X) \not\equiv v(X) \equiv \sin X \quad \text{but} \quad V(X) = v(x) \quad \text{for all} \quad x = X^2$$

Quiz

$$\mathbf{V}(\mathbf{X}, t) = \mathbf{V}(\mathbf{X}(\mathbf{x}, t), t) \equiv \mathbf{v}(\mathbf{x}, t)$$


# trajectory of a particle



trajectory:  $\vec{x}(\vec{x}_0, t)$

velocity:  $\vec{v} = \frac{d}{dt} \vec{x}(\vec{x}_0, t)$


## trajectory of a particle



trajectory:  $\vec{x}(\vec{x}_0, t)$

velocity:  $\vec{v} = \frac{d}{dt} \vec{x}(\vec{x}_0, t)$

# trajectory of a particle



trajectory:  $\vec{x}(\vec{x}_0, t)$

velocity:  $\vec{v} = \frac{d}{dt} \vec{x}(\vec{x}_0, t)$



# Velocity

---

Parametric derivative

$$\text{velocity} = \frac{d}{dt} \mathbf{x}(\mathbf{X}_0, t)$$





# Velocity

---

Parametric derivative

$$\text{velocity} = \frac{d}{dt} \mathbf{x}(\mathbf{X}_0, t)$$

Definition

$$\mathbf{V}(\mathbf{X}, t) = \frac{\partial \mathbf{x}}{\partial t}$$



# Velocity

---

Parametric derivative

$$\text{velocity} = \frac{d}{dt} \mathbf{x}(\mathbf{X}_0, t)$$

Definition

$$\mathbf{V}(\mathbf{X}, t) = \frac{\partial \mathbf{x}}{\partial t} = \frac{\partial \mathbf{u}}{\partial t}$$



# Velocity

---

Parametric derivative

$$\text{velocity} = \frac{d}{dt} \mathbf{x}(\mathbf{X}_0, t)$$

Definition

$$\mathbf{V}(\mathbf{X}, t) = \frac{\partial \mathbf{x}}{\partial t} = \frac{\partial \mathbf{u}}{\partial t}$$

Parallel transport

$$\mathbf{V}(\mathbf{X}, t) = \mathbf{V}(\mathbf{X}(\mathbf{x}, t), t) \equiv \mathbf{v}(\mathbf{x}, t)$$



# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$



# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$

## Chain rule calculation

$$\frac{\partial \mathbf{V}}{\partial t} \equiv \frac{\partial}{\partial t} \mathbf{v}(\mathbf{x}(\mathbf{X}, t), t)$$





# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$

## Chain rule calculation

$$\frac{\partial \mathbf{V}}{\partial t} \equiv \frac{\partial}{\partial t} \mathbf{v}(x_1(\mathbf{X}, t), x_2(\mathbf{X}, t), x_3(\mathbf{X}, t), t)$$



# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$

## Chain rule calculation

$$\frac{\partial \mathbf{V}}{\partial t} \equiv \frac{\partial}{\partial t} \mathbf{v}(x_1(\mathbf{X}, t), x_2(\mathbf{X}, t), x_3(\mathbf{X}, t), t) = \frac{\partial \mathbf{v}}{\partial x_j} \frac{\partial x_j}{\partial t} + \frac{\partial \mathbf{v}}{\partial t}$$



# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$

## Chain rule calculation

$$\frac{\partial \mathbf{V}}{\partial t} \equiv \frac{\partial}{\partial t} \mathbf{v}(x_1(\mathbf{X}, t), x_2(\mathbf{X}, t), x_3(\mathbf{X}, t), t) = \frac{\partial \mathbf{v}}{\partial x_j} V_j(\mathbf{X}, t) + \frac{\partial \mathbf{v}}{\partial t}$$



# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$

## Chain rule calculation

$$\frac{\partial \mathbf{V}}{\partial t} \equiv \frac{\partial}{\partial t} \mathbf{v}(x_1(\mathbf{X}, t), x_2(\mathbf{X}, t), x_3(\mathbf{X}, t), t) = \frac{\partial \mathbf{v}}{\partial x_j} v_j(\mathbf{x}, t) + \frac{\partial \mathbf{v}}{\partial t}$$



# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$

## Chain rule calculation

$$\frac{\partial \mathbf{V}}{\partial t} \equiv \frac{\partial}{\partial t} \mathbf{v}(x_1(\mathbf{X}, t), x_2(\mathbf{X}, t), x_3(\mathbf{X}, t), t) = \frac{\partial \mathbf{v}}{\partial x_j} v_j(\mathbf{x}, t) + \frac{\partial \mathbf{v}}{\partial t} = \mathbf{a}(\mathbf{x}, t)$$

No L-field needed!





# Acceleration

---

## Definition

$$\mathbf{A}(\mathbf{X}, t) = \frac{\partial \mathbf{V}}{\partial t} \mapsto \mathbf{a}(\mathbf{x}, t)$$

## Chain rule calculation

$$\frac{\partial \mathbf{V}}{\partial t} \equiv \frac{\partial}{\partial t} \mathbf{v}(x_1(\mathbf{X}, t), x_2(\mathbf{X}, t), x_3(\mathbf{X}, t), t) = \frac{\partial \mathbf{v}}{\partial x_j} v_j(\mathbf{x}, t) + \frac{\partial \mathbf{v}}{\partial t} = \mathbf{a}(\mathbf{x}, t)$$

$$\mathbf{a} = \frac{\partial \mathbf{v}}{\partial t} + \frac{\partial \mathbf{v}}{\partial x_j} v_j$$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = X_1 + at^2$$

$$x_2 = X_2 + bX_2t + ct^2$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = X_1 + at^2$$

$$x_2 = X_2 + bX_2t + ct^2$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(X_1, X_2, t) = 2at$$

$$V_2(X_1, X_2, t) = bX_2 + 2ct$$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = X_1 + at^2$$

$$x_2 = X_2 + bX_2t + ct^2$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(X_1, X_2, t) = 2at$$

$$V_2(X_1, X_2, t) = bX_2 + 2ct$$

Euler description

$$v_1(x_1, x_2, t) = 2at$$

$$v_2(x_1, x_2, t) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = X_1 + at^2$$

$$x_2 = X_2 + bX_2t + ct^2$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(X_1, X_2, t) = 2at$$

$$V_2(X_1, X_2, t) = bX_2 + 2ct$$

Euler description

$$v_1(x_1, x_2, t) = 2at$$

$$v_2(x_1, x_2, t) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$

Numerical example:  $X_1 = X_2 = 0, t = 3$





## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = X_1 + at^2$$

$$x_2 = X_2 + bX_2t + ct^2$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(X_1, X_2, t) = bX_2 + 2ct$$

Euler description

$$v_1(x_1, x_2, t) = 2at$$

$$v_2(x_1, x_2, t) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$

Numerical example:  $X_1 = X_2 = 0, t = 3$





## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = X_1 + at^2$$

$$x_2 = X_2 + bX_2t + ct^2$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(0, 0, 3) = 6c$$

Euler description

$$v_1(x_1, x_2, t) = 2at$$

$$v_2(x_1, x_2, t) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$

Numerical example:  $X_1 = X_2 = 0, t = 3$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = 9a$$

$$x_2 = X_2 + bX_2t + ct^2$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(0, 0, 3) = 6c$$

Euler description

$$v_1(x_1, x_2, t) = 2at$$

$$v_2(x_1, x_2, t) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$

Numerical example:  $X_1 = X_2 = 0, t = 3$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = 9a$$

$$x_2 = 9c$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(0, 0, 3) = 6c$$

Euler description

$$v_1(x_1, x_2, t) = 2at$$

$$v_2(x_1, x_2, t) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$

Numerical example:  $X_1 = X_2 = 0, t = 3$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = 9a$$

$$x_2 = 9c$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(0, 0, 3) = 6c$$

Euler description

$$v_1(9a, 9c, 3) = 2at$$

$$v_2(9a, 9c, 3) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$

Numerical example:  $X_1 = X_2 = 0, t = 3$



## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = 9a$$

$$x_2 = 9c$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(0, 0, 3) = 6c$$

Euler description

$$v_1(9a, 9c, 3) = 6a$$

$$v_2(9a, 9c, 3) = b \frac{x_2 - ct^2}{1 + 3b} + 6c$$

Numerical example:  $X_1 = X_2 = 0$ ,  $t = 3$





## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = 9a$$

$$x_2 = 9c$$

Inverse

$$X_1 = x_1 - at^2$$

$$X_2 = \frac{x_2 - ct^2}{1 + bt}$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(0, 0, 3) = 6c$$

Euler description

$$v_1(9a, 9c, 3) = 6a$$

$$v_2(9a, 9c, 3) = b \frac{9c - 9c}{1 + 3b} + 6c$$

Numerical example:  $X_1 = X_2 = 0, t = 3$





## Example 1: Domain mapping (1/2)

---

Mapping

$$x_1 = 9a$$

$$x_2 = 9c$$

Inverse

$$X_1 = 0$$

$$X_2 = 0$$

Lagrange description

$$V_1(0, 0, 3) = 6a$$

$$V_2(0, 0, 3) = 6c$$


Euler description

$$v_1(9a, 9c, 3) = 6a$$



$$v_2(9a, 9c, 3) = 6c$$

Numerical example:  $X_1 = X_2 = 0, t = 3$



For a fixed  $t_0$ :



For a fixed  $t_0$ :



For a fixed  $t_0$ :



Same values, different functions, different derivatives!



## Example 1: Domain mapping (2/2)

---

$$V_1(X_1, X_2, t) = 2at$$

$$V_2(X_1, X_2, t) = bX_2 + 2ct$$



## Example 1: Domain mapping (2/2)

---

$$V_1(X_1, X_2, t) = 2at \quad A_1(X_1, X_2, t) = 2a = a_1(x_1, x_2, t)$$

$$V_2(X_1, X_2, t) = bX_2 + 2ct \quad A_2(X_1, X_2, t) = 2c = a_2(x_1, x_2, t)$$





## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$ ,  $A_2 = a_2 = 2c$

Euler field

$$v_1(x_1, x_2, t) = 2at$$

$$v_2(x_1, x_2, t) = b \frac{x_2 - ct^2}{1 + bt} + 2ct$$

## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$ ,  $A_2 = a_2 = 2c$

Euler field

$$\begin{aligned} v_1(x_1, x_2, t) &= 2at & a_1(x_1, x_2, t) &= \frac{\partial v_1}{\partial t} + \frac{\partial v_1}{\partial x_j} v_j \\ v_2(x_1, x_2, t) &= b \frac{x_2 - ct^2}{1 + bt} + 2ct & a_2(x_1, x_2, t) &= \frac{\partial v_2}{\partial t} + \frac{\partial v_2}{\partial x_j} v_j \end{aligned}$$

## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$  (checks),  $A_2 = a_2 = 2c$

Euler field

$$\begin{aligned} v_1(x_1, x_2, t) &= 2at & a_1(x_1, x_2, t) &= \frac{\partial v_1}{\partial t} + \frac{\partial v_1}{\partial x_j} v_j = 2a \\ v_2(x_1, x_2, t) &= b \frac{x_2 - ct^2}{1 + bt} + 2ct & a_2(x_1, x_2, t) &= \frac{\partial v_2}{\partial t} + \frac{\partial v_2}{\partial x_j} v_j \end{aligned}$$



## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$  (checks),  $A_2 = a_2 = 2c$

Euler field

$$\begin{aligned} v_1(x_1, x_2, t) &= 2at & a_1(x_1, x_2, t) &= \frac{\partial v_1}{\partial t} + \frac{\partial v_1}{\partial x_j} v_j = 2a \\ v_2(x_1, x_2, t) &= b \frac{x_2 - ct^2}{1 + bt} + 2ct & a_2(x_1, x_2, t) &= \frac{\partial v_2}{\partial t} + \frac{\partial v_2}{\partial x_j} v_j \end{aligned}$$

---

$$\frac{\partial v_2}{\partial t} =$$

+

$$\frac{\partial v_2}{\partial x_j} v_j =$$

## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$  (checks),  $A_2 = a_2 = 2c$

Euler field

$$\begin{aligned} v_1(x_1, x_2, t) &= 2at & a_1(x_1, x_2, t) &= \frac{\partial v_1}{\partial t} + \frac{\partial v_1}{\partial x_j} v_j = 2a \\ v_2(x_1, x_2, t) &= b \frac{x_2 - ct^2}{1 + bt} + 2ct & a_2(x_1, x_2, t) &= \frac{\partial v_2}{\partial t} + \frac{\partial v_2}{\partial x_j} v_j \end{aligned}$$

---

$$\frac{\partial v_2}{\partial t} = b \frac{-2ct(1 + bt) - (x_2 - ct^2)b}{(1 + bt)^2} + 2c$$

+

$$\frac{\partial v_2}{\partial x_j} v_j =$$



## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$  (checks),  $A_2 = a_2 = 2c$

Euler field

$$\begin{aligned} v_1(x_1, x_2, t) &= 2at & a_1(x_1, x_2, t) &= \frac{\partial v_1}{\partial t} + \frac{\partial v_1}{\partial x_j} v_j = 2a \\ v_2(x_1, x_2, t) &= b \frac{x_2 - ct^2}{1 + bt} + 2ct & a_2(x_1, x_2, t) &= \frac{\partial v_2}{\partial t} + \frac{\partial v_2}{\partial x_j} v_j \end{aligned}$$

---

$$\frac{\partial v_2}{\partial t} = b \frac{-2ct(1 + bt) - (x_2 - ct^2)b}{(1 + bt)^2} + 2c$$

+

$$\frac{\partial v_2}{\partial x_j} v_j = \frac{b}{1 + bt} \left( b \frac{x_2 - ct^2}{1 + bt} + 2ct \right)$$



## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$  (checks),  $A_2 = a_2 = 2c$

Euler field

$$\begin{aligned}
 v_1(x_1, x_2, t) &= 2at & a_1(x_1, x_2, t) &= \frac{\partial v_1}{\partial t} + \frac{\partial v_1}{\partial x_j} v_j = 2a \\
 v_2(x_1, x_2, t) &= b \frac{x_2 - ct^2}{1 + bt} + 2ct & a_2(x_1, x_2, t) &= \frac{\partial v_2}{\partial t} + \frac{\partial v_2}{\partial x_j} v_j
 \end{aligned}$$

$$\left. \begin{aligned}
 \frac{\partial v_2}{\partial t} &= b \frac{-2ct(1 + bt) - (x_2 - ct^2)b}{(1 + bt)^2} + 2c \\
 + \\
 \frac{\partial v_2}{\partial x_j} v_j &= \frac{b}{1 + bt} \left( b \frac{x_2 - ct^2}{1 + bt} + 2ct \right)
 \end{aligned} \right\} = 2c$$

## Example 1: Domain mapping (2/2)

---

stored result:  $A_1 = a_1 = 2a$  (checks),  $A_2 = a_2 = 2c$  (checks)

Euler field

$$\begin{aligned}
 v_1(x_1, x_2, t) &= 2at & a_1(x_1, x_2, t) &= \frac{\partial v_1}{\partial t} + \frac{\partial v_1}{\partial x_j} v_j = 2a \\
 v_2(x_1, x_2, t) &= b \frac{x_2 - ct^2}{1 + bt} + 2ct & a_2(x_1, x_2, t) &= \frac{\partial v_2}{\partial t} + \frac{\partial v_2}{\partial x_j} v_j = 2c
 \end{aligned}$$

$$\left. \begin{aligned}
 \frac{\partial v_2}{\partial t} &= b \frac{-2ct(1 + bt) - (x_2 - ct^2)b}{(1 + bt)^2} + 2c \\
 + \\
 \frac{\partial v_2}{\partial x_j} v_j &= \frac{b}{1 + bt} \left( b \frac{x_2 - ct^2}{1 + bt} + 2ct \right)
 \end{aligned} \right\} = 2c$$